

MULTIMEDIA



UNIVERSITY

STUDENT ID NO

--	--	--	--	--	--	--	--	--	--

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 3, 2017/2018

DIM5068 – MATHEMATICAL TECHNIQUES 2

(For DIT students only)

4th JUNE 2018

9.00 a.m – 11.00 a.m

(2 Hours)

INSTRUCTIONS TO STUDENT

1. This question paper consists of 2 pages with 4 questions excluded the cover page and Appendix. Key formulae are given in the Appendix.
2. Answer ALL questions.
3. Write your answers in the answer booklet provided.
4. All necessary working steps must be shown.

Question 1

a. Differentiate the following functions with respect to x by using **Chain Rule**.

i) $f(x) = \cos(x^3 - 5x^2 - 4x)$. (5 marks)

ii) $y = 2 \ln(3x^2 + 4)$ (6 marks)

b. Differentiate $x^3y + 2y^2 - 7x^2 = x^2y$ by using **implicit differentiation**.

(7 marks)

c. Given $f(x) = 4x^3 + 6x^2 - 24x - 12$

i) Find the intervals on which the function is increasing and decreasing (5 marks)

ii) Identify the function's local extreme values (2 marks)

[TOTAL 25 MARKS]

Question 2


a. Find/evaluate the following integrals:

i) $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (4x^2 + \sin x) dx$ (5 marks)

ii) $\int (4x^3)(2x^4 - 2)^6 dx$ [Hint: Use **Substitution Rule**] (5 marks)

iii) $\int_0^2 \frac{x}{\sqrt{x^2}} dx$ [Hint: Use **Substitution Rule**] (7 marks)

b. Find the area of the region R bounded by $y = 2x^2 - 4x + 6$, $y = -x^2 + 2x + 1$, $x = 1$ and $x = 2$. (8 marks)



[TOTAL 25 MARKS]

Continued...

Question 3

a. Solve the following differential equation by using the **Separable Method**.

i) $\sqrt{2+4x^3} dy = y^4 x^2 dx$. (Hint: **Solve for y**) (8 marks)

ii) $3\frac{dy}{dx} = 6x^2 y + 3y$. (Hint: **No need to solve for y**) (4 marks)

b. Given the differential equation $4\frac{dy}{dx} + 8y = 4e^{-2x}$

i) Identify the $p(x)$ and $q(x)$. (3 marks)

ii) Calculate the integrating factor, μ . (1.5 marks)

iii) Find y given $\mu y = \int \mu q(x) dx$. (3.5marks)

[TOTAL 20 MARKS]

Question 4

a. Given $\vec{a} = \langle -3, 2, 1 \rangle$ and $\vec{b} = \langle 2, 0, -2 \rangle$, find

i) $3\vec{a} + 4\vec{b}$ (3 marks)

ii) $|3\vec{a} + 4\vec{b}|$ (2 marks)

b. Find the angle between $\vec{u} = 4\vec{i} - 3\vec{j}$ and $\vec{v} = 2\vec{i} + 5\vec{j}$. (5 marks)

c. Determine whether $\vec{v} = 3\vec{i} - 2\vec{j}$ and $\vec{w} = 4\vec{i} + 6\vec{j}$ are parallel, orthogonal, or neither. (3 marks)

d. Given the vertices of a triangle $X = (0, -2, 0)$, $Y = (4, 1, -2)$, and $Z = (5, 3, 1)$.

i) Determine \vec{XY} and \vec{XZ} . (2 marks)

ii) Calculate the cross product of \vec{XY} and \vec{XZ} . (3 marks)

iii) Compute the total area of the triangle. Correct your answer to 2 decimal places. (3 marks)

e. If a line passing through the points $(1, 3, 2)$ and $(-4, 2, 0)$, compute the

i) parametric equations of the line. (4 marks)

ii) symmetric equations of the line. (3 marks)

f. Find an **equation of the plane** that passes through the point $(1, 4, -5)$ and is perpendicular to $3\vec{i} + 2\vec{j} - 2\vec{k}$. Leave your answer in the form of $ax + by + cz = d$. (2 marks)

[TOTAL 30 MARKS]

End of page.

APPENDIX

Derivatives: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

Differentiation Rules

General Formulae

$$1. \frac{d}{dx}[f(x)g(x)] = f(x)g'(x) + g(x)f'(x) \quad 2. \frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2}$$

$$3. \frac{d}{dx}(x^n) = nx^{n-1} \quad 4. \frac{d}{dx}[f(u)] = \frac{dy}{du} \cdot \frac{du}{dx}$$

Exponential and Logarithmic Functions

$$1. \frac{d}{dx}(e^x) = e^x \quad 2. \frac{d}{dx}(a^x) = a^x \ln a$$

$$3. \frac{d}{dx}(\ln x) = \frac{1}{x} \quad 4. \frac{d}{dx}(\log_a x) = \frac{1}{x \ln a}$$

Trigonometric Functions

$$1. \frac{d}{dx}(\sin x) = \cos x \quad 2. \frac{d}{dx}(\cos x) = -\sin x$$

$$3. \frac{d}{dx}(\tan x) = \sec^2 x \quad 4. \frac{d}{dx}(\csc x) = -\csc x \cot x$$

$$5. \frac{d}{dx}(\sec x) = \sec x \tan x \quad 6. \frac{d}{dx}(\cot x) = -\csc^2 x$$

Table of Integrals

$$1. \int u \, dv = uv - \int v \, du \quad 2. \int u^n \, du = \frac{u^{n+1}}{n+1} + C, \quad n \neq -1$$

$$3. \int \frac{du}{u} = \ln|u| + C \quad 4. \int e^u \, du = e^u + C$$

$$5. \int \sin u \, du = -\cos u + C \quad 6. \int \cos u \, du = \sin u + C$$

$$7. \int \sec^2 u \, du = \tan u + C \quad 8. \int \csc^2 u \, du = -\cot u + C$$

$$9. \int \sec u \tan u \, du = \sec u + C \quad 10. \int \csc u \cot u \, du = -\csc u + C$$

Application of Integrals:

Areas between Curve, $A = \int_a^b [f(x) - g(x)] \, dx$

Differential Equations

Linear Differential Equations

$$\frac{dy}{dx} + p(x)y = q(x) \quad \Rightarrow \quad \mu y = \int \mu q(x) dx, \text{ where } \mu = e^{\int p(x) dx}$$

Constant Coefficient of Homogeneous Equations

$$\text{Roots of Auxiliary Equation, } r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

General Solutions to the Auxiliary Equation:

$$\begin{array}{ll} 2 \text{ Real \& Unequal Roots } (b^2 - 4ac > 0) & y = c_1 e^{r_1 x} + c_2 e^{r_2 x} \\ \text{Repeated Roots } (b^2 - 4ac = 0) & y = c_1 e^{rx} + c_2 x e^{rx} \\ 2 \text{ Complex Roots } (b^2 - 4ac < 0) & y = e^{ax} (c_1 \cos bx + c_2 \sin bx) \end{array}$$

Constant Coefficient of Non-Homogeneous Equations

$$y = y_c + y_p \text{ [} y_c : \text{complementary solution, } y_p : \text{particular solution]}$$

Vector

Length of Vector

$$\text{The length of the vector } \mathbf{a} = \langle a_1, a_2, a_3 \rangle \text{ is } |\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}.$$

Dot Product

If θ is the angle between the vector $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$, then

$$\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2 + a_3 b_3 = |\mathbf{a}| |\mathbf{b}| \cos \theta$$

Cross Product

If θ is the angle between the vector $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$, then

$$\mathbf{a} \times \mathbf{b} = \langle a_2 b_3 - a_3 b_2, a_3 b_1 - a_1 b_3, a_1 b_2 - a_2 b_1 \rangle$$

$$|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$$

Area for parallelogram PQRS

$$= \left| \vec{PQ} \times \vec{PR} \right|$$

Area for triangle PQR

$$= \frac{1}{2} \left| \vec{PQ} \times \vec{PR} \right|$$

Equation of Lines

Vector equation: $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$

Parametric equations: $x = x_0 + at \quad y = y_0 + bt \quad z = z_0 + ct$

$$\text{Symmetric equation: } \frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}$$

Equation of Planes

Vector equation: $\mathbf{n} \cdot \mathbf{r} = \mathbf{n} \cdot \mathbf{r}_0$

Scalar equations: $a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$

Linear equation: $ax + by + cz + d = 0$

$$\text{Angle between Two Planes: } \theta = \cos^{-1} \left(\frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{|\mathbf{n}_1| |\mathbf{n}_2|} \right)$$